

ALLERGY/ANAPHYLAXIS ACTION PLAN

Student Name _____ D.O.B. _____ Grade _____
 Address _____ Phone Number _____
 Name of Doctor _____ Preferred Hospital _____

Student

Photo

ALLERGY: (check appropriate) To be completed by Doctor

- Foods (list):
 Medications (list):
 Latex: Circle: Type I (anaphylaxis) Type IV (contact dermatitis)
 Stinging Insects (list):

RECOGNITION AND TREATMENT

Chart to be completed by Physician ONLY		Give CHECKED Medication	
If food ingested or contact with allergen occurs:		EpiPen®	Antihistamine
No symptoms noted	I <input type="checkbox"/> Observe for other symptoms		
Minor symptoms - Such as itchy mouth, few hives, mild itch, mild nausea/discomfort			
Mouth	Itching, tingling, or swelling of lips, tongue, mouth		
Skin	Hives, itchy rash, swelling of the face or extremities		
Gut+	Nausea, abdominal cramps, vomiting, diarrhea		
Throat+	Tightening of throat, hoarseness, hacking cough		
Lung+	Shortness of breath, repetitive coughing, wheezing		
Heart+	Thready pulse, low BP, fainting, pale, blueness		
Neuro+	Disorientation, dizziness, loss of conscience		
If reaction is progressing (several of the above areas affected), The severity of symptoms can quickly change. +Potentially life-threatening.			

DOSAGE:

Epinephrine: Inject into outer thigh **EpiPen® 0.3 mg** OR **EpiPen® Jr. 0.15 mg** (see reverse for instructions)

Antihistamine: Benadryl _____ .mg To be given by mouth *only if able to swallow.*

This child has received instruction in the proper use of the EpiPen®. It is my professional opinion that this student **SHOULD** be allowed to carry and use the EpiPen® independently. The child knows when to request antihistamine and has been advised to inform a responsible adult if the EpiPen® is self-administered.

It is my professional opinion that this student **SHOULD NOT** carry the EpiPen®.

EMERGENCY CALLS

1. Call 911. State that an allergic reaction has been treated, and additional epinephrine may be needed.
2. Call parents/guardian to notify of reaction, treatment and student's health status.
3. Treat for shock. Prepare to do CPR.
4. Accompany student to ER if no parent/guardians are available.

PHYSICIAN'S SIGNATURE. _____ **DATE** _____

Side 2: To Be Completed by Parent/Guardian, Student and School

Allergy/Anaphylaxis Action Plan (continued) Student Name _____

Parent/Guardian AUTHORIZATIONS

- I want this allergy plan implemented for my child; I want my child to carry the EpiPen® and I agree to release the school district and school personnel from all claims of liability if my child suffers any adverse reactions from self-administration of EpiPen®.
- I want this plan implemented for my child and I do not want my child to self-administer EpiPen®.
- It is recommended that backup medication be stored with the school nurse in case a student forgets or loses the EpiPen® and/or antihistamine. The school district is not responsible or liable if backup medication is not provided to the school nurse and the student is without working medication when medication is needed.

Your signature gives permission for the nurse to contact and receive additional information from your health care provider regarding the allergic condition(s) and the prescribed medication.

Parent/Guardian Signature: _____ **Date:** _____

Student Agreement:

- I have been trained in the use of my EpiPen® and allergy medication and understand the signs and symptoms for which they are given;
- I agree to carry my EpiPen® with me at all times;
- I will notify a responsible adult (teacher, nurse, coach, noon duty, etc.) IMMEDIATELY when auto-injector EpiPen® (epinephrine) is used;
- I will not share my medication with other students or leave my EpiPen® unattended;
- I will not use my allergy medications for any other use than what it is prescribed for.

Student Signature _____

- Back-up medication is stored at school Yes No
-

DIRECTIONS FOR EPIPEN® USE

1. Pull off gray activation cap.
2. Hold black tip to outer thigh (apply to thigh only).
3. Press hard into outer thigh until auto-injector mechanism functions. Hold in place for 10 seconds.
4. Massage the injection site for 10 seconds.
Other: _____
5. Once EpiPen® is used, call 911/EMS. Take the used EpiPen® to the emergency room with you.

STAFF MEMBERS TRAINED

NAME	TITLE	ROOM	TRAINED BY

EMERGENCY CONTACTS

		NAME	HOME #	WORK #	CELL #
Parent/Guardian					
Parent/Guardian					
Other:					